

# PFIZER BIVALENT

## Individuals 6 months through 4 years of age:

|   |   |
|---|---|
| Unvaccinated individuals:   | Three doses of Pfizer Bivalent are administered. The first two doses are given 3 weeks apart. The third dose is given at least 8 weeks after the second dose.   |
| Individuals who have received one dose of a monovalent Pfizer Vaccine:    | Two doses of Pfizer Bivalent are administered. The first dose is given 3 weeks after the monovalent Pfizer and the second dose is given at least 8 weeks later. |
| Individuals who have received two doses of a monovalent Pfizer Vaccine:   | A single dose of Pfizer Bivalent is administered at least 8 weeks after the last monovalent Pfizer vaccine.   |
| Individuals who have received three doses of a monovalent Pfizer Vaccine: | A single dose of Pfizer Bivalent is administered at least 2 months after the last monovalent Pfizer vaccine.  |

## Individuals 5 years of age and older:

|   |   |
|---|---|
| Unvaccinated individuals:   | A single dose of Pfizer Bivalent is administered.   |
| Individuals who have received one or more doses of a monovalent COVID-19 Vaccine:                                 | A single dose of Pfizer Bivalent is administered at least 2 months after the last monovalent COVID-19 vaccine.  |
| Individuals 65 years of age and older who have received one dose of a bivalent COVID-19 vaccine:                  | A single dose of Pfizer Bivalent may be administered at least 4 months after one dose of bivalent COVID-19 vaccine. This is an optional dose; any adult 65 years of age or older who has received a single bivalent dose is up-to-date.             |
| Immunocompromised individuals 5 years of age and older who have received one dose of a bivalent COVID-19 vaccine: | An additional dose of Pfizer Bivalent may be administered at least 2 months following their one dose of bivalent COVID-19 vaccine. This is an optional dose; immunocompromised individuals who have received a single bivalent dose are up-to-date. |

# MODERNA BIVALENT

## Individuals 6 months through 5 years of age:

|  |   |
|--|---|
| Unvaccinated individuals:  | Two doses of Moderna Bivalent are administered. The second dose is given 1 month after the first.           |
| Individuals who have received one dose of a monovalent Moderna Vaccine:  | A single dose of Moderna Bivalent is administered 1 month after the last dose of Moderna vaccine.           |
| Individuals who have received two doses of a monovalent Moderna Vaccine: | A single dose of Moderna Bivalent is administered at least 2 months after the last dose of Moderna vaccine. |

## Individuals 6 years of age and older:

|  |  |
|--|--|
| Unvaccinated individuals:  | A single dose of Moderna Bivalent is administered.   |
| Individuals who have received one or more doses of a monovalent COVID-19 Vaccine:                | A single dose of Moderna Bivalent is administered at least 2 months after any monovalent COVID-19 vaccine.             |
| Individuals 65 years of age and older who have received one dose of a bivalent COVID-19 vaccine: | A single dose of Moderna Bivalent may be administered at least 4 months after one dose of a bivalent COVID-19 vaccine. |

## Immunocompromised individuals 6 years of age and older:

|   |   |
|---|---|
| Immunocompromised individuals 6 years and older | A single, additional age-appropriate dose of Moderna Bivalent may be administered at least 2 months following the initial dose of a bivalent COVID-19 vaccine |
|---|---|